arXiv:1110.2548v2 [hep-ph] 3 Feb 2012

Preprint typeset in JHEP style - PAPER VERSION

Deep inelastic scattering of baryons in a modified soft wall model

Nelson R. F. Braga

Instituto de Física, Universidade Federal do Rio de Janeiro, Caixa Postal 68528,
RJ 21941-972 – Brazil

Alfredo Vega

Departamento de Física y Centro Científico Tecnológico de Valparaíso
(CCTVal), Universidad Técnica Federico Santa María, Casilla 110-V,
Valparaíso, Chile

E-mails: `braga@if.ufrj.br`, `alfredo.vega@usm.cl`

ABSTRACT: We calculate the structure functions for unpolarized deep inelastic scattering of baryons using an AdS/QCD soft wall model that considers a dressed mass for the bulk fermionic fields. Considering the regime of large Bjorken parameter x, we compare the results for the proton structure function F_2 with experimental results.

KEYWORDS: Gauge-gravity correspondence, AdS-CFT Correspondence, Deep Inelastic Scattering.

Contents

1. Introduction 1

2. Deep Inelastic Scattering and Structure Functions 2

3. DIS in the soft wall model with fermionic dressed bulk mass 4

4. Analysis of the results 10

1. Introduction

Our present knowledge of strong interactions indicates that QCD (Quantum Chromodynamics) is the quantum field theory that models the physics of this fundamental interaction. However, many important properties of the strongly interacting particles, the hadrons, can not be described using only perturbative QCD. This happens because the QCD coupling is large at low energies. Thus, in order to calculate static properties of hadrons, or to describe their structure, one needs additional tools.

In the last years, many interesting models to study non perturbative aspects of hadronic physics were developed based on the idea of gauge/string duality. The main source of inspiration for these kind of models was the discovery of the AdS/CFT correspondence[1, 2, 3] that is an exact duality between string theory in certain ten dimensional geometries and supersymmetric $SU(N)$ gauge theories with large N on the corresponding boundary. In particular, string theory in $AdS_5 \times S^5$ space is dual to a four dimensional gauge theory.

In the AdS/CFT correspondence, the gauge theory is conformal. The idea of breaking this conformal invariance by an Infrared (IR) cut off, represented in the dual AdS geometry by a geometrical cut off in the radial coordinate of the space, was introduced in [4] as a tool to reproduce the high energy scaling of hadronic scattering amplitudes for processes at fixed angles. This scaling had been found in QCD a long time before [5, 6] but the corresponding string theory description was lacking for a very long time.

This approach of considering a maximum radial size of AdS space as the dual of an infrared cut off in the gauge theory was then used in [7, 8] to calculate the mass spectrum of glueballs. In these articles, boundary conditions were imposed on fields living in an AdS slice and the corresponding normal modes were associated with hadronic states. This kind of model, later called AdS/QCD hard wall, was then applied to other hadrons, as, for example, in [9].

The hadronic masses calculated using the hard wall model do not exhibit the observed approximate linear relation between mass squared and spin (and radial excitation number), corresponding to the so called linear Regge trajectories. Rather, the hard wall masses show asymptotically quadratic trajectories. This motivated a different AdS/QCD approach consisting of a background involving AdS space plus a field that acts effectively as a smooth infrared cut off [10]. This, so called soft wall model, leads to linear Regge trajectories for mesons (see also [11] for glueballs).

For fermions, the soft wall model, as originally formulated[10], does not lead to a discrete spectrum because the dilaton introduced in the action is factorized out in the equations of motion. This situation can be changed, following alternative approaches in order to reproduce Regge behavior for baryons. One example can be found in [12, 13], where the authors consider models in asymptotically AdS space including a warp factor in the metric. Other possibility, studied in [14, 15, 16], considers a z dependent (or dressed) mass for the fermionic modes propagating in AdS space. This last approach has been considered in [15] to study nucleon form factors, and in [17] to obtain some generalized parton distributions (GPD) for nucleons. Here we consider baryons in an AdS/QCD soft wall model with dressed mass in order to describe deep inelastic scattering (DIS).

The study of deep inelastic scattering (DIS) using the AdS/CFT correspondence appeared first in [18]. Then other authors considered the description of DIS using various AdS/QCD models like, for example, in [19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29]. In [19] the soft wall AdS/QCD model was considered for scalar hadrons and a hybrid model involving a soft wall cut off for the photons and a hard wall cut off for the fermions was also discussed. However, the calculation of baryonic structure functions using a soft wall cut off was still missing. We will use the modified soft wall model with dressed mass, studied in [15, 16] to calculate the DIS structure functions for baryons. Then we compare our F_2 with experimental results for the proton in the regime of large values of the Bjorken parameter x where the supergravity approximation that we use is more reliable.

In section **2** we present a briefly review of DIS and hadronic structure functions. In section **3** we show our AdS/QCD calculation of the structure functions. Then in section **4** we analyze our results, discussing the possible choices of the parameters of the model, plotting the structure function F_2 for some kinematical regimes and comparing with experimental results.

2. Deep Inelastic Scattering and Structure Functions

Deep inelastic scattering (DIS) is a process where a highly energetic lepton, an electron in general, interacts with a hadronic target through the exchange of a virtual



Figure 1: Diagram of a deep inelastic scattering.

photon, as show in the diagram of figure **1**. The momenta of the photon and of the initial hadron are respectively q^μ and P^μ. After the interaction there is a final hadronic state represented by X with momentum P_X^μ. The experimental measurement of the inclusive cross section of DIS corresponds to detecting the final lepton, thus determining the momentum transfer q^μ, but not the final hadronic state X. That means, summing over all possible final hadronic states X. One usually parametrizes DIS using as dynamical variables the photon virtuality q^2 and the Bjorken parameter

$$x \equiv -q^2/2P\cdot q\,. \tag{2.1}$$

Deep inelastic scattering, in the strict sense, corresponds to the limit $q^2 \to \infty$, with x fixed.

The inclusive cross section for DIS can be calculated from the hadronic tensor that is defined as

$$W^{\mu\nu} = \frac{1}{8\pi}\sum_s \int d^4y\, e^{iq\cdot y}\langle P,s|\, \Big[J^\mu(y), J^\nu(0)\Big]\,|P,s\rangle\,, \tag{2.2}$$

where $J^\mu(y)$ is the electromagnetic hadronic current and s the spin of the initial hadron. For an unpolarized scattering (spin independent) this tensor can be decomposed in terms of the structure functions $F_1(x,q^2)$ and $F_2(x,q^2)$ as [30]

$$W^{\mu\nu} = F_1(x,q^2)\Big(\eta^{\mu\nu} - \frac{q^\mu q^\nu}{q^2}\Big) + \frac{2x}{q^2}F_2(x,q^2)\Big(P^\mu + \frac{q^\mu}{2x}\Big)\Big(P^\nu + \frac{q^\nu}{2x}\Big)\,. \tag{2.3}$$

Considering only final states with just one baryon with mass M_X, we can introduce a basis of such one particle states and write the hadronic tensor as

$$\begin{aligned} W^{\mu\nu} = {} & \frac{1}{8\pi}\sum_{s,s_X}\sum_{M_X}\int \frac{d^4P_X}{2\pi^3)}\theta(P_X^0)\delta\Big(P_X^2 + M_X^2\Big)(2\pi)^4\delta^4(P+q-P_X) \\ & \times \langle P,s|J^\nu(0)|P_X,s_X\rangle\,\langle P_X,\, s_X|J^\mu(0)|P,\, s\rangle \end{aligned}$$

$$= \frac{1}{4} \sum_{s, s_X} \sum_{M_X} \delta\Big(M_X^2 + (P+q)^2 \Big) \langle P, s | J^\nu(0) | P+q, s_X \rangle \, \langle P+q, \, s_X | J^\mu(0) | P, \, s \rangle \; . \tag{2.4}$$

So, in order to calculate the hadronic structure functions one needs to find the matrix elements of the current and also the spectrum of hadronic masses M_X . We will present in the next section the calculation of these quantities using the modified soft wall model.

3. DIS in the soft wall model with fermionic dressed bulk mass

In the soft wall model, a four dimensional gauge theory is represented by a gravity dual consisting of fields living in anti-de Sitter space AdS_5, whose metric can be written as

$$ds^2 \equiv g_{mn}\, dy^m dy^n \; = \; \frac{R^2}{z^2} (dz^2 + \eta_{\mu\nu} dy^\mu dy^\nu) \; = \; \frac{R^2}{z^2} (dz^2 - dt^2 + (d\vec{y})^2) \; , \tag{3.1}$$

with the presence of an additional background of the form $e^{-\kappa^2 z^2}$. This factor plays the role of an infrared cut off, with the energy parameter κ representing the corresponding scale. The general form of an action integral corresponding to a given Lagrangian density $\mathcal{L}$ is

$$I \; = \; \int dz d^4y \, \sqrt{-g} \; e^{-\kappa^2 z^2} \, \mathcal{L} \; . \tag{3.2}$$

For the case of scalar and vector fields, this kind of action integral leads to dual four dimensional theories with infrared cut off and discrete mass spectra. For fermions one must do some modification in order to find a dual theory with an infrared cut off. This happens because using a standard fermionic Lagrangian in the action above the background $e^{-\kappa^2 z^2}$ factors out of the equations of motion. So, the fermions are not affected by the energy scale κ and get no discrete mass spectrum. This problem can be solved introducing a z dependent mass, as studied in [15, 16]. Following this approach, the appropriate action that describes the dynamics of the fermionic and gauge fields and their interaction is

$$\begin{aligned} I \; = \; \int dz d^4y \, \sqrt{-g} \, e^{-\kappa^2 z^2} \, \Big[& -\frac{1}{4} F_{mn} F^{mn} \; + \; \frac{i}{2} \bar{\Psi} \epsilon_a^m \Gamma^a \mathcal{D}_m \Psi \\ & - \; \frac{i}{2} (\mathcal{D}_m \Psi)^\dagger \Gamma^0 \epsilon_a^m \Gamma^a \Psi \; - \; \bar{\Psi} \Big(\mu + V_F(z) \Big) \Psi \Big] \; , \end{aligned} \tag{3.3}$$

where the field strength is $F^{mn} = \partial^m A^n - \partial^n A^m$ and $\epsilon_a^m = \delta_a^m \frac{z}{R}$ and the covariant derivative is:

$$\mathcal{D}_m = \partial_m - \frac{1}{8}\omega_m^{ab}\,[\Gamma_a, \Gamma_b] - i g_{_V} A_m\,, \tag{3.4}$$

with: $\omega_m^{ab} = -\frac{1}{z}(\delta_z^a \delta_m^b - \delta_z^b \delta_m^a)$. The coupling $g_{_V}$ will be associated with the electric charge e and $\Gamma^a = (\gamma^\mu, -i\gamma^5)$ are the Dirac matrices.

In order to produce a discrete spectrum with the Regge behavior for the fermions we introduced the effective fermionic potential $V_F(z) = \kappa^2 z^2/R$ depending on the z coordinate (associated with the boundary energy scale), to dress the fermionic bulk mass.

The physical motivation for this procedure is the following. On one hand, according to the AdS/CFT dictionary, the mass of a supergravity bulk field is related to the dimension of the corresponding boundary operator. On the other hand, in general the dimensions of quantum operators receive anomalous contributions, depending on the energy scale. Since the energy scale of the boundary theory is holographycally related to the localization in the z coordinate, the possibility of anomalous contributions to the dimension of the operators can be translated into z dependent masses for the dual bulk modes [14, 31]. This kind of procedure makes it possible to introduce an important ingredient of QCD that is not considered in most of the AdS / QCD models. Other related works that consider masses varying in the bulk can be found in, for example, [32, 33, 34, 35].

For the gauge fields it is convenient [18] to impose the gauge condition

$$\partial_\mu A^\mu \,+\, z e^{\kappa^2 z^2} \partial_z \Big(e^{-\kappa^2 z^2} \frac{1}{z} A_z \Big) \,=\, 0\,. \tag{3.5}$$

Note that from now on we use the notation of raising and lowering the four dimensional indices with the Minkowski metric: $A^\mu \equiv \eta^{\mu\nu} A_\nu$ and $\Box \equiv \eta^{\mu\nu}\partial_\mu\partial_\nu$. With the gauge choice (3.5) the equations of motion that emerge from the action (3.3) are

$$\begin{aligned} &\Box A^\mu \,+\, z e^{\kappa^2 z^2} \partial_z \Big(e^{-\kappa^2 z^2} \frac{1}{z} \partial_z A^\mu \Big) \,=\, 0 \\ &\Box A_z \,-\, \partial_z \Big(\partial_\mu A^\mu \Big) \,=\, 0\,. \end{aligned} \tag{3.6}$$

We impose the condition that the boundary value of the gauge field represents a virtual photon with polarization η^μ and space-like momentum q^μ

$$A_\mu(z,y)|_{z\to 0} \,=\, \eta_\mu\, e^{iq\cdot y}\,, \tag{3.7}$$

The corresponding solutions are

$$\begin{aligned} A_\mu(z,y) &= \eta_\mu\, e^{iq\cdot y}\, \kappa^2\; \Gamma(1+\frac{q^2}{4\kappa^2})\; z^2\; \mathcal{U}(1+\frac{q^2}{4\kappa^2}; 2; \kappa^2 z^2) \\ A_z(z,y) &= \frac{i}{2}\,\eta\cdot q\, e^{iq\cdot y}\; \Gamma(1+\frac{q^2}{4\kappa^2})\; z\; \mathcal{U}(1+\frac{q^2}{4\kappa^2}; 1; \kappa^2 z^2)\,, \end{aligned} \tag{3.8}$$

where $\mathcal{U}(a;b;w)$ are the confluent hypergeometric functions of the second kind.

Now, regarding the fermionic fields, it is convenient to do the following field redefinition

$$\Psi(y,z) = e^{+\kappa^2 z^2/2}\psi(y,z). \tag{3.9}$$

in such a way that the equations of motion take the form

$$\Big[i\,\not{\partial} + \gamma^5 \partial_z - \frac{2}{z}\gamma^5 - \frac{1}{z}\Big(m + \kappa^2 z^2\Big)\Big]\psi(y,z) = 0\,, \tag{3.10}$$

where $\not{\partial} = \gamma^\mu\,\partial_\mu$. The quantity μ, the bulk fermion mass was replaced by the dimensionless parameter $m = \mu R$. In order to solve the equation of motion, we decompose the fermionic field into chiral components

$$\psi(y,z) = \psi_L(y,z) + \psi_R(y,z)\,, \quad \psi_{L/R} = \frac{1 \mp \gamma^5}{2}\psi\,. \tag{3.11}$$

with $(\gamma^5)^2 = 1$.

We will consider solutions of the form

$$\psi_{L/R}(y,z) = e^{iP\cdot y}\frac{1}{2}\Big(1 \mp \gamma^5\Big)u_s(P)\,\frac{z^2}{R^2} f_{L/R}(z)\,, \tag{3.12}$$

that contain the plane wave factor and the four component spinor $u_s(P)$ corresponding to a four dimensional free fermion with momentum P^μ and spin s. Then, the z dependent parts of chiral components of the fields: $f_{L/R}(z)$ must satisfy

$$\Big[-\partial_z^2 + \kappa^4 z^2 + 2\kappa^2\Big(m \mp \frac{1}{2}\Big) + \frac{m(m \pm 1)}{z^2}\Big] f_{L/R}(z) = -P^2\, f_{L/R}(z)\,. \tag{3.13}$$

Following the usual prescription of gauge/gravity dualities, normalizable solutions for fields in the gravity side are dual to states in the boundary four dimensional theory. Equations (3.13) have normalizable solutions only when $-P^2$, the four dimensional mass, has the discrete values

$$-P_n^2 = M_n^2 = 4\kappa^2\Big(n + m + \frac{1}{2}\Big)\,, \tag{3.14}$$

with $n = 0, 1, 2, ...,$. The corresponding discrete set of normalizable solutions $f^n_{L/R}(z)$ are

$$f_L^n(z) = \sqrt{\frac{2\Gamma(n+1)}{\Gamma(n+m+3/2)}}\;\kappa^{m+3/2}\;z^{m+1}\;e^{-\kappa^2 z^2/2}\;L_n^{m+1/2}(\kappa^2 z^2)\,, \tag{3.15}$$

$$f_R^n(z) = \sqrt{\frac{2\Gamma(n+1)}{\Gamma(n+m+1/2)}}\;\kappa^{m+1/2}\;z^{m}\;e^{-\kappa^2 z^2/2}\;L_n^{m-1/2}(\kappa^2 z^2) \tag{3.16}$$

with normalization condition $\int\limits_0^\infty dz\, f^m_{L/R}(z) f^n_{L/R}(z) = \delta_{mn}$.

We want to describe processes where the initial state is a proton that absorbs a virtual photon transforming into a final state corresponding to an excited hadronic state of spin 1/2. So we consider the interaction action

$$S_{int}[i,X] = g_{_V} \int dz d^4y \sqrt{-g} e^{-\kappa^2 z^2}\, \frac{z}{R}\, A_m \bar{\Psi}_X\, \gamma^m\, \Psi_i\,. \tag{3.17}$$

where Ψ_i represents the initial proton, that we take as the state with lowest mass level, corresponding to $n = 0$. So the initial momentum $P_i \equiv p$ satisfies $-p^2 = M_0^2 = 4\kappa^2(m + 1/2)$. The fermionic field Ψ_X represents a final state with (higher) mass M_X and momentum $P_X = p + q$ satisfying $-P_X^2 = M_X^2 = 4\kappa^2(n_X + m + 1/2)$, where we are representing as n_X the integer associated with the excitation level of the final state. The corresponding solutions have the form

$$\begin{aligned} \Psi_i &= e^{ip\cdot y} e^{+\kappa^2 z^2/2}\, \frac{z^2}{R^2} \Big[\Big(\frac{1-\gamma^5}{2}\Big) u_{s_i}(p)\, f^0_L(z) \,+\, \Big(\frac{1+\gamma^5}{2}\Big) u_{s_i}(p)\, f^0_R(z) \Big] \\ \Psi_X &= e^{iP_X\cdot y} e^{+\kappa^2 z^2/2}\, \frac{z^2}{R^2} \Big[\Big(\frac{1-\gamma^5}{2}\Big) u_{s_X}(P_X)\, f^{n_X}_L(z) \,+\, \Big(\frac{1+\gamma^5}{2}\Big) u_{s_X}(P_X)\, f^{n_X}_R(z) \Big]\,, \end{aligned} \tag{3.18}$$

where s_i and s_X are the spins of the initial and final fermionic states.

In order to find out the structure functions for the hadrons we have to calculate the interaction action (3.17) with the solutions for the fields. We can simplify the calculations by considering, as it was done in [18, 19], that we are probing the hadron with a particular photon with polarization η^μ satisfying $\eta \cdot q = 0$. For this situation the z component of the gauge field does not contribute and, substituting the solutions (3.6) and (3.18) in (3.17) the interaction action, for this initial and final states (i, X) takes the on shell form

$$\begin{aligned} S_{int}[i,X] = \frac{g_{_V}}{2}\, (2\pi)^4 \delta^4(P_X - p - q) \eta_\mu \Big[& \bar{u}_{s_X}(P_X) \gamma^\mu \Big(\frac{1-\gamma^5}{2}\Big) u_{s_i}(p) \mathcal{I}_L(n_x) \\ & + \bar{u}_{s_X}(P_X) \gamma^\mu \Big(\frac{1+\gamma^5}{2}\Big) u_{s_i}(p) \mathcal{I}_R(n_x) \Big]\,, \end{aligned} \tag{3.19}$$

where $\mathcal{I}_L(n_x)$ and $\mathcal{I}_R(n_x)$ are integrals involving the two fermionic solutions with different chiralities. They have a similar structure and can be written, in terms of the variable $w \equiv \kappa^2 z^2$, in the general form:

$$\mathcal{I}(\bar{m}, n_X) = C(\bar{m}, n_X)\, \Gamma(1 + \frac{q^2}{4\kappa^2}) \int_0^\infty dw w^{\bar{m}-1} e^{-w} \mathcal{U}(1 + \frac{q^2}{4\kappa^2}; 2; w)\, L^{\bar{m}-2}_{n_X}(w)\,, \tag{3.20}$$

where

$$C(\bar{m}, n_X) = \sqrt{\frac{4\Gamma(n_X+1)}{\Gamma(\bar{m}-1)\Gamma(n_X+\bar{m}-1)}} \,. \tag{3.21}$$

The integrals $\mathcal{I}_L(n_x)$ and $\mathcal{I}_R(n_x)$ correspond to $\mathcal{I}(\bar{m}, n_X)$ with $\bar{m} = m + 5/2$ and $\bar{m} = m + 3/2$ respectively. Performing the integral (3.20) we get

$$\mathcal{I}(\bar{m}, n_X) = \frac{q^2 \Gamma\left(\bar{m}\right) \sqrt{\frac{\Gamma(\bar{m}-1)\Gamma(n_X+\bar{m}-1)}{\Gamma(n_X+1)}} \Gamma\left(\frac{q^2}{4\kappa^2} + n_X\right)}{2\kappa^2 \Gamma\left(\bar{m}-1\right) \Gamma\left(\frac{q^2}{4\kappa^2} + n_X + \bar{m}\right)} \tag{3.22}$$

Now that we calculated the action in the soft wall background for photons interacting with fermions with dressed mass, we connect this result with the four dimensional boundary theory using a proposal similar to the one used in refs. [18, 19]. In these references the matrix element of the fermionic electromagnetic current on the boundary four dimensional theory was considered to be equal to the corresponding bulk interaction action. Here we will take a different point of view and consider that bulk/boundary duality implies that these quantities are proportional, rather than necessarily equal. So we assume the relations:

$$\begin{aligned} \eta_\mu \langle P_X | \tilde{J}^\mu(q) | P \rangle \; &= (2\pi)^4 \, \delta^4(P_X - P - q) \, \eta_\mu \, \langle P + q | J^\mu(0) | P \rangle \; = \; \mathcal{K}_{_{eff}} \, S_{int}[i, X] \\ \eta_\mu \langle P | \tilde{J}^\mu(q) | P_X \rangle \; &= (2\pi)^4 \, \delta^4(P_X - P - q) \, \eta_\mu \, \langle P | J^\mu(0) | P + q \rangle \; = \; \mathcal{K}_{_{eff}} \, S_{int}[X, i] \,, \end{aligned} \tag{3.23}$$

where $\mathcal{K}_{_{eff}}$ plays the role of a bulk/boundary effective factor that phenomenologically adjust the bulk supergravity quantities to the boundary observed ones. Following this prescription, the hadronic tensor in eq. (2.4), contracted with the photon polarization η, can then be written in terms of our interaction action of eq. (3.19) as

$$\begin{aligned} \eta_\mu \eta_\nu W^{\mu\nu} \; &= \; \frac{1}{4} \sum_{M_X} \delta(M_X^2 + (p+q)^2) \, \frac{g_{eff}^2}{4} \times \\ \sum_{s_i} \sum_{s_X} \; & \Big\{ \mathcal{I}_L^2 \mathcal{I}_R^2 \Big(\bar{u}_{s_X} \gamma^\mu \Gamma^{(-)} u_{s_i} \bar{u}_{s_i} \gamma^\nu \Gamma^{(+)} u_{s_X} + \bar{u}_{s_X} \gamma^\mu \Gamma^{(+)} u_{s_i} \bar{u}_{s_i} \gamma^\nu \Gamma^{(-)} u_{s_X} \Big) \\ + & \mathcal{I}_L^2 \bar{u}_{s_X} \gamma^\mu \Gamma^{(-)} u_{s_i} \bar{u}_{s_i} \gamma^\nu \Gamma^{(-)} u_{s_X} \; + \mathcal{I}_R^2 \bar{u}_{s_X} \gamma^\mu \Gamma^{(+)} u_{s_i} \bar{u}_{s_i} \gamma^\nu \Gamma^{(+)} u_{s_X} \Big\} \end{aligned} \tag{3.24}$$

where we defined $g_{eff} \equiv \mathcal{K}_{_{eff}} \, g_{_V}$ and $\Gamma^{(\pm)} \equiv \frac{1 \pm \gamma^5}{2}$ and omitted the dependence of the spinors on the momenta. Note that we are summing over the final spins and averaging over the initial ones. Using the property

$$\sum_s (u_s)_\alpha(p) (\bar{u}_s)_\beta(p) = (\gamma^\mu p_\mu + m)_{\alpha\beta} \,,$$

satisfied by both the initial and final state (on shell) spinors, with the corresponding masses and momenta, we find

$$\eta_\mu \eta_\nu W^{\mu\nu} = \frac{1}{4} \sum_{M_X} \delta(M_X^2 + (p+q)^2) \; g_{eff}^2 \Big\{ - \mathcal{I}_L^2(n_x) \mathcal{I}_R^2(n_x) M_X M_0 \, \eta \cdot \eta \\ + \Big(\mathcal{I}_L^2(n_x) + \mathcal{I}_R^2(n_x) \Big) \Big((p \cdot \eta)^2 - \frac{1}{2} (p^2 + p \cdot q) \, \eta \cdot \eta \Big) \Big\} \qquad (3.25)$$

The sum over the final states X can be approximated by an integral over a continuum of states as it was done in in refs. [18, 19]. In the present case we this corresponds to replacing the sum over X of the delta functions by the factor: $\frac{1}{4\kappa^2}$. That means:

$$\eta_\mu \eta_\nu W^{\mu\nu} \approx \frac{g_{eff}^2}{16\kappa^2} \Big\{ - \mathcal{I}_L^2(n_x) \mathcal{I}_R^2(n_x) M_X M_0 \, \eta \cdot \eta \\ + \Big(\mathcal{I}_L^2(n_x) + \mathcal{I}_R^2(n_x) \Big) \Big((p \cdot \eta)^2 - \frac{1}{2} (p^2 + p \cdot q) \, \eta \cdot \eta \Big) \Big\} \qquad (3.26)$$

For the particular photon that we are considering, with $\eta \cdot q = 0$ we get from eq. (2.3)

$$\eta_\mu \eta_\nu W^{\mu\nu} \; = \; \eta^2 F_1 \; + \; \frac{2x}{q^2} (\eta \cdot p)^2 \, F_2 \qquad (3.27)$$

Comparing this with our expression for the hadronic tensor eq. (3.26) we find our results for the fermionic structure functions

$$F_1 = \frac{g_{eff}^2}{16\kappa^2} \Big\{ \Big(\mathcal{I}_L^2(n_x) + \mathcal{I}_R^2(n_x) \Big) \Big(\frac{M_0^2}{2} + \frac{q^2}{4x} \Big) \\ - \mathcal{I}_L(n_x) \mathcal{I}_R(n_x) M_0 \sqrt{M_0^2 + \frac{q^2(1-x)}{x}} \Big\} \\ F_2 = \frac{g_{eff}^2}{32\kappa^2} \Big(\mathcal{I}_L^2(n_x) + \mathcal{I}_R^2(n_x) \Big) \frac{q^2}{x} \qquad (3.28)$$

The excitation level n_x of the final state is not an independent variable. It can be expressed in terms the DIS variables q^2 and x. Using eqs. (2.1) and (3.14) one finds

$$- (p+q)^2 = M_0^2 + q^2 \Big[\frac{1}{x} - 1 \Big] = M_X^2 = 4\kappa^2 \Big(n_X + m + \frac{1}{2} \Big) , \qquad (3.29)$$

so

$$n_X \; = \; \frac{q^2}{4\kappa^2} \Big[\frac{1}{x} - 1 \Big] , \qquad (3.30)$$



Figure 2: F_2 as a function of q^2, where q^2 is in units of $(\text{GeV})^2$. The plot consider $x = 0.85$. The higher curve in for $m = 0.6$, the middle one for $m = 0.7$ and the lower is plotted with $m = 0.8$. Dots correspond to experimental data [36, 37].

4. Analysis of the results

In order to calculate the structure functions and analyze their dependence on x and q^2, first we have to fix the parameters of the model. The effective coupling $g_{eff} \equiv \mathcal{K}_{eff}\, g_V$ contains the electric charge g_V, that satisfies $g_V^2 = 1/137$ but multiplied by the yet undetermined parameter $\mathcal{K}_{eff}$ inserted in the model in order to phenomenologically adjust the relation between bulk and boundary quantities and essentially fix the size the structure functions. We found that the choice $g_{eff}^2 = 0.66$ leads to structure functions F_2 with values compatible with experimental data. So we used this value in the calculations.

The infrared energy scale κ is related with the Regge slope. In fact, in the present holographic model the Regge slope is $4\kappa^2$. We will choose $4\kappa^2 = 0.9\, GeV^2$ that was found in ref. [12, 14] to give a good adjustment for the trajectory of nucleons. So we use $\kappa = 0.474 GeV$. The other parameter of the model: m, is considered as a free parameter, associated, as we discussed before, with the anomalous dimension of the operator that creates the baryonic states. We used this free parameter to look for the best fit of the shape of the structure function F_2 to the experimental data. It is important to remark that the supergravity approximation used in the model is, in principle, valid only for large values of x. This happens because, as discussed in [18], for low values of x string theory corrections would become relevant. So we analyzed the region $0.8 < x < 1.0$. We show in Fig. 2 the structure functions found with the holographic model for $x = 0.85$ compared with the corresponding experimental data that appear in PDG [36, 37] considering the values of m that led to the best fits, that means, m close to 0.7.

In order to check if the dependence of F_2 on q^2 has a similar form for other values of x in the range considered, we show in Fig. 3 this structure function for



Figure 3: F_2 as a function of q^2. The plot on the left consider $x = 0.8$ and on the right $x = 0.9$. In both plots the higher curve in for $m = 0.6$, the middle one for $m = 0.7$ and the lower is plotted with $m = 0.8$.

$x = 0.8$ and $x = 0.9$. In both cases we included plots for three different choices of the parameter m. We see that F_2 decreases with q^2 in a way that is compatible with the experimental results shown in figure (16.7) of [36] or in [37] for $x = 0.85$.

Now, let's consider the dependence of F_2 on the Bjorken parameter x. There are not many experimental results for this structure function for x close to one. But one finds in ref. [38] an interesting investigation of F_2 at large x, for low values of q^2, using some parametrizations obtained from experimental data. In particular, they show results for the F_2 as a function of x for the cases of $q^2 = 4$ GeV^2 and 9 GeV^2. Their results show a decrease in this structure function when we increase x in the interval $0.9 < x < 1$.

We show in Fig. 4 the structure function F_2 obtained using our holographic model with z dependent mass in the range $0.8 < x < 1.0$, for these two values of q^2 analyzed in [38], using values for m around 0.7. The order of magnitude of our results was adjusted by the choice of the effective coupling g_{eff} to be consistent with the experimental values, so they are of the same order of those found in [38], however, in contrast to the results of [38], we found an increase in the structure functions when $x \to 1$. So, our model for DIS of baryons does not give a good description for the dependence of F_2 on x at low q^2. This may be a consequence of the fact that in this simple model the final hadronic states have just one baryon with spin 1/2. In a general non elastic process the final state may include more than one hadron and also hadrons with higher spins. A description of such kind of process with multiple hadrons and also higher spins is out of the scope of the present kind of model. Nevertheless it is interesting to see that the model is able to reproduce, by adjusting the free parameters, the experimental dependence of F_2 on q^2.

Finally, it is interesting to consider the high q limit of the structure functions eq.(3.28). Using the property



Figure 4: F_2 v/s x. The plot at the left consider $q^2 = 4GeV^2$ and the other is for $q^2 = 8GeV^2$. In both plots the higher curve in for $m = 0.6$, the middle one for $m = 0.7$ and the lower is plotted with $m = 0.8$.

$$\frac{\Gamma(a+y)}{\Gamma(b+y)} = \left(\frac{1}{y}\right)^{b-a}\Big(1+O(y^{-1})\Big); \qquad y \to \infty\,, \tag{4.1}$$

we can expand the integrals $I_{L,R}$ defined by eq. (3.22) and that appear in eq.(3.28). This way we find properties shared with other AdS/QCD models at dominant order in q, like

$$F_2 = 2F_1 \tag{4.2}$$

that implies that for $x \to 1$ we approach the Callan-Gross relation. We also find

$$F_2 \sim \left(\frac{q^2}{4\kappa^2}\right)^{-m+\frac{1}{2}} x^{m+\frac{5}{2}}(1-x)^{m+\frac{5}{2}}. \tag{4.3}$$



Figure 5: F_2 v/s x. The plot at the left consider $q^2 = 30\ GeV^2$ and the other is for $q^2 = 1000\ GeV^2$. In both plots the higher curve in for $m = 0.6$, the middle one for $m = 0.7$ and the lower is plotted with $m = 0.8$. It can be seen that at high q the structure functions decrease for $x \to 1$.

Notice that when $m = \tau - 3/2$ our model reproduces results found in previous works that consider DIS in holographic models [18, 19]. Additionally, at high values for q^2, we find a decreasing behavior for the structure function when x is close to one, as can be seen in Fig. 5.

Acknowledgments: We would like to thank Carlos Alfonso Ballon Bayona for important discussions. N.B. is partially supported by CNPq and Capes (Brazil) and A.V is supported by Fondecyt (Chile) under Grant No. 3100028. A.V. is grateful for the hospitality of the Instituto de Física of Universidade Federal do Rio de Janeiro, where this work started.

References

[1] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)]. [arXiv:hep-th/9711200].

[2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998). [arXiv:hep-th/9802109].

[3] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998). [arXiv:hep-th/9802150].

[4] J. Polchinski and M. J. Strassler, Phys. Rev. Lett. **88**, 031601 (2002) [arXiv:hep-th/0109174].

[5] V. A. Matveev, R. M. Muradian and A. N. Tavkhelidze, Lett. Nuovo Cim. **7**, 719 (1973).

[6] S. J. Brodsky and G. R. Farrar, Phys. Rev. Lett. **31**, 1153 (1973); Phys. Rev. D **11**, 1309 (1975).

[7] H. Boschi-Filho and N. R. F. Braga, Eur. Phys. J. C **32**, 529 (2004) [arXiv:hep-th/0209080].

[8] H. Boschi-Filho and N. R. F. Braga, JHEP **0305**, 009 (2003) [arXiv:hep-th/0212207].

[9] G. F. de Teramond and S. J. Brodsky, Phys. Rev. Lett. **94**, 201601 (2005) [arXiv:hep-th/0501022].

[10] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. D **74**, 015005 (2006) [arXiv:hep-ph/0602229].

[11] P. Colangelo, F. De Fazio, F. Jugeau and S. Nicotri, Phys. Lett. B **652**, 73 (2007) [arXiv:hep-ph/0703316].

[12] H. Forkel, M. Beyer and T. Frederico, JHEP **0707**, 077 (2007) [arXiv:0705.1857 [hep-ph]].

[13] W. de Paula, T. Frederico, H. Forkel and M. Beyer, Phys. Rev. D **79**, 075019 (2009) [arXiv:0806.3830 [hep-ph]].

[14] A. Vega, I. Schmidt, Phys. Rev. **D79**, 055003 (2009). [arXiv:0811.4638 [hep-ph]].

[15] Z. Abidin and C. E. Carlson, Phys. Rev. D **79**, 115003 (2009) [arXiv:0903.4818 [hep-ph]].

[16] T. Gutsche, V. E. Lyubovitskij, I. Schmidt and A. Vega, arXiv:1108.0346 [hep-ph].

[17] A. Vega, I. Schmidt, T. Gutsche, V. E. Lyubovitskij, Phys. Rev. **D83**, 036001 (2011). [arXiv:1010.2815 [hep-ph]].

[18] J. Polchinski and M. J. Strassler, JHEP **0305**, 012 (2003) [arXiv:hep-th/0209211].

[19] C. A. Ballon Bayona, H. Boschi-Filho and N. R. F. Braga, JHEP **0803**, 064 (2008) [arXiv:0711.0221 [hep-th]].

[20] C. A. Ballon Bayona, H. Boschi-Filho and N. R. F. Braga, JHEP **0810**, 088 (2008) [arXiv:0712.3530 [hep-th]].

[21] L. Cornalba and M. S. Costa, Phys. Rev. D **78**, 096010 (2008) [arXiv:0804.1562 [hep-ph]].

[22] B. Pire, C. Roiesnel, L. Szymanowski and S. Wallon, Phys. Lett. B **670**, 84 (2008) [arXiv:0805.4346 [hep-ph]].

[23] J. L. Albacete, Y. V. Kovchegov and A. Taliotis, JHEP **0807**, 074 (2008) [arXiv:0806.1484 [hep-th]].

[24] J. H. Gao and B. W. Xiao, Phys. Rev. D **80** (2009) 015025 [arXiv:0904.2870 [hep-ph]].

[25] Y. Hatta, T. Ueda and B. W. Xiao, JHEP **0908** (2009) 007 [arXiv:0905.2493 [hep-ph]].

[26] C. A. Ballon Bayona, H. Boschi-Filho and N. R. F. Braga, JHEP **0809**, 114 (2008) [arXiv:0807.1917 [hep-th]].

[27] L. Cornalba, M. S. Costa and J. Penedones, JHEP **1003**, 133 (2010) [arXiv:0911.0043 [hep-th]].

[28] Y. Hatta, E. Iancu and A. H. Mueller, JHEP **0801**, 063 (2008) [arXiv:0710.5297 [hep-th]].

[29] Y. Y. Bu and J. M. Yang, Phys. Rev. D **84**, 106004 (2011) [arXiv:1109.4283 [hep-th]].

[30] For a review see: A. V. Manohar, arXiv:hep-ph/9204208.

[31] A. Cherman, T. D. Cohen, E. S. Werbos, Phys. Rev. **C79**, 045203 (2009). [arXiv:0804.1096 [hep-ph]].

[32] H. Forkel, E. Klempt, Phys. Lett. **B679**, 77-80 (2009). [arXiv:0810.2959 [hep-ph]].

[33] H. Forkel, Phys. Lett. **B694**, 252-257 (2010). [arXiv:1007.4341 [hep-ph]].

[34] A. Vega, I. Schmidt, Phys. Rev. **D82**, 115023 (2010). [arXiv:1005.3000 [hep-ph]].

[35] A. Vega, I. Schmidt, Phys. Rev. **D84**, 017701 (2011). [arXiv:1104.4365 [hep-ph]].

[36] K. Nakamura *et al.* [Particle Data Group Collaboration], J. Phys. G **G37**, 075021 (2010).

[37] L. W. Whitlow, E. M. Riordan, S. Dasu, S. Rock and A. Bodek, Phys. Lett. B **282**, 475 (1992).

[38] S. P. Malace *et al.* [Jefferson Lab E00-115 Collaboration], Phys. Rev. **C80**, 035207 (2009). [arXiv:0905.2374 [nucl-ex]].